

Heinz Body Hemolytic Anemia: A Step by Step Illustration

Basil M. KAHWASH¹; Geok Chin TAN^{2,3}, Samir B. KAHWASH²

¹ Indiana University School of Medicine, Department of Internal Medicine, Indianapolis IN, USA

² Nationwide Children's Hospital, Department of Pathology and Laboratory Medicine, Columbus OH, USA

³ National University of Malaysia, Department of Pathology, Kuala Lumpur, MALAYSIA

The images demonstrate the morphologic changes of RBCs in peripheral blood smear of a patient with G6PD deficiency. Figure 1 includes three RBCs (demarcated by numbered arrows) that represent the successive stages of this phenomenon. Stage 1: Precipitated hemoglobin appears as a dense round zone within the RBC (arrow 1). Stage 2: Precipitated hemoglobin was removed upon passage through a reticuloendothelial system-rich tissue, leaving a blister-like vacuole (arrow 2). Stage 3: An RBC transforms into a bite cell (arrow 3). The transformation from stage 1 to stage 2 and 3 is expedited as RBCs pass through the narrow lumens of capillaries.




Figure 1. Morphological stages of formation of bite cells: Precipitated haemoglobin (Heinz body) appears as a dense spot at the periphery of RBC (arrow 1), to a RBC with a blister-like vacuole (arrow 2) and eventually to a bite cell in which the precipitated haemoglobin was removed (arrow 3).




Figure 2. RBCs travel through tight luminal space of a capillary (arrow 4).

Figure 2 contains a capillary from a stained touch imprint that illustrates the tight luminal space that RBCs must travel through (arrow 4). Precipitated hemoglobin appears as blue inclusions at periphery of RBC's (Figure 3- arrows) stained with supravital or Brilliant Cresyl Blue.

Heinz body was first described by Robert Heinz in cases with hemolytic anemia in 1890.¹ The underlying pathophysiology involves denaturation and precipitation of hemoglobin due to an inability of G6PD-deficient RBCs to clear oxygenated free radicals. RBCs deformed by precipitated hemoglobin lack the flexibility needed for tight navigation, particularly while passing through sinusoids of the spleen.^{2,3}




Figure 3. Precipitated haemoglobin appears as blue inclusion at the periphery of RBCs in supravital stain (multiple arrows)

REFERENCES

1. Heinz R. Morphologische Veränderungen des roten Blutkörperchens durch Gifte. [Virchows] Archiv für pathologische Anatomie und Physiologie und für klinische Medizin, Berlin 122: 112-116, 1890.
2. Fujita T, Kashimura M, Adachi K. Scanning electron microscopy (SEM) studies of the spleen--normal and pathological. Scan Electron Microsc (Pt 1): 435-444, 1982.

Correspondence:

Geok CHIN TAN
Department of Pathology
National University of Malaysia
Jalan Yaacob Latif, Bandar Tun Razak
56000 KUALA LUMPUR / MALAYSIA

Tel: 603-91455362
e-mail: tangc@ppukm.ukm.edu.my